

Review

The Vienna consensus: report of an expert meeting on the development of ART laboratory performance indicators



ESHRE Special Interest Group of Embryology and Alpha Scientists in Reproductive Medicine ^{a,b,*}

^a European Society of Human Reproduction and Embryology, Meerstraat 60, B-1852 Grimbergen, Belgium

^b ALPHA Scientists in Reproductive Medicine, 19 Mayıs Mah. 19 Mayıs Cad. Nova Baran Center No:4 34360 Sisli, Istanbul, Turkey

KEY MESSAGE

This proceedings report presents 19 Indicators, including 12 Key Performance Indicators (KPIs), 5 Performance Indicators (PIs), and 2 Reference Indicators (RIs) from an international workshop supported by the European Society of Human Reproduction and Embryology (ESHRE) and Alpha Scientists in Reproduction (Alpha), designed to establish consensus on definitions and recommended values for the assisted reproductive technology (ART) laboratory.

ABSTRACT

This proceedings report presents the outcomes from an international workshop supported by the European Society of Human Reproduction and Embryology (ESHRE) and Alpha Scientists in Reproductive Medicine, designed to establish consensus on definitions and recommended values for Indicators for the assisted reproductive technology (ART) laboratory. Minimum performance-level values ('competency') and aspirational ('benchmark') values were recommended for a total of 19 Indicators, including 12 Key Performance Indicators (KPIs), five Performance Indicators (PIs), and two Reference Indicators (RIs).

© 2017 The Author(s). Published by Elsevier Ltd on behalf of Reproductive Healthcare Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Performance Indicators (PIs) are objective measures for evaluating critical healthcare domains (patient safety, effectiveness, equity, patient-centeredness, timeliness and efficiency) (Kohn et al., 2000). In the setting of a clinical laboratory, quality indicators are necessary

for systematically monitoring and evaluating the laboratory's contribution to patient care (ISO-15189:2012) and they represent an important element within the quality management system (QMS) (ESHRE Guideline Group on Good Practice in IVF Labs et al, 2016; Mortimer and Mortimer, 2015). Currently, there are no established PIs for assisted reproductive technology (ART) laboratories, and there is very little published evidence on the topic.

This consensus document, which has not been subjected to independent peer review by the editors of *Reproductive BioMedicine Online*, is being published simultaneously by *Reproductive BioMedicine Online* and *Human Reproduction Open*.

* Corresponding authors.

E-mail addresses: coticchio.biogenesi@grupposandonato.it (G Coticchio); zsolt.peter.nagy@gmail.com (ZP Nagy).
<https://doi.org/10.1016/j.rbmo.2017.06.015>

1472-6483/© 2017 The Author(s). Published by Elsevier Ltd on behalf of Reproductive Healthcare Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Any PI should be reliable and robust, and routine data collection for the indicator should be straightforward. Furthermore, the biological or technical process to be monitored should be defined, and relevant qualifiers, confounders and endpoints should be identified. Key Performance Indicators (KPIs) are Indicators deemed essential for evaluating the introduction of a technique or process; establishing minimum standards for proficiency; monitoring ongoing performance within a QMS (for internal quality control (IQC), external quality assurance (EQA)); benchmarking and quality improvement. In general, the results of a series of KPIs will provide an adequate overview of the most important steps in the IVF laboratory process (Salinas et al., 2010).

The aim of the consensus meeting and report was to establish KPIs for ART laboratories for use in monitoring 'fresh' IVF and ICSI cycles and provide the basis for several of the quantitative performance criteria needed to create competency profiles for clinical embryologists. More specifically, the purpose was to achieve an international consensus regarding: (i) a minimum list of IVF laboratory indicators and KPIs that in the future can be further extended and/or revised; (ii) specific definitions for these indicators (including necessary case inclusion/exclusion criteria; and calculation formulae); and (iii) recommended values for each KPI (minimum 'competency' limit; and 'aspirational goal' benchmark).

Based on the information presented here, each laboratory should develop its own set of KPIs founded on laboratory organization and processes, and develop a systematic, transparent, and consistent approach to data collection and analysis and calculation of KPIs (ESHRE Guideline Group on Good Practice in IVF Labs et al., 2016; Mayer et al., 2003; Mortimer and Mortimer, 2015; Salinas et al., 2010).

Methodology

This report is the result of a 2-day consensus meeting of expert professionals (participants are listed in Table 1). As a starting point for the discussion at the meeting, two surveys were organized to collect information on indicators used in IVF laboratories. The first, the 'Alpha survey', was sent to national and international societies of ART laboratory directors and clinical embryologists, and to the members of the European Society of Human Reproduction and Embryology (ESHRE) committee of national representatives. Eighteen responses were received out of 34 sent, with opinions from Australia, Austria, Belgium, Bulgaria, Canada, Croatia, France, Germany, Ireland, Italy, Japan, Slovenia, Sweden, South Africa, Turkey, UK, and USA, and the results of this survey informed the expert panel on minimum expected, or competence, values (i.e. values that any laboratory should be able to achieve), and aspirational, or benchmark, values (i.e. values that can be employed as a best practice goal), for a range of quality indicators. Where possible, responses were based on standardized information (national collected data, or large datasets), but in most countries such data are not available. Another survey, the 'ESHRE survey', provided information on current practice (How many KPIs are measured, frequency of measurement, characteristics of a reference population for KPIs) and the degree of importance of some indicators. This survey was sent to 2413 members of the ESHRE Special Interest Group (SIG) of Embryology, and 384 responses were received. In addition, where relevant, published data were collected from a literature search and summarized, although for most indicators,

Table 1 – Consensus workshop participants and contributors.

Participant/contributor name	Affiliation
Susanna Apter	Fertilitetscentrum Stockholm, Sweden
Basak Balaban	American Hospital of Istanbul, Turkey
Alison Campbell ^a	CARE Fertility Group, UK
Jim Catt	Optimal IVF, Melbourne, Australia
Giovanni Coticchio	Biogenesi, Monza, Italy
Maria José de los Santos ^a	IVI Valencia, Valencia, Spain
Sophie Debrock ^a	Leuven University Fertility Centre, Leuven, Belgium
Thomas Ebner ^a	Kepler University, Linz, Austria
Stephen Harbottle	Cambridge IVF, UK
Ciara Hughes	Rotunda IVF, Dublin, Ireland
Ronny Janssens	Centre for Reproductive Medicine, Brussels, Belgium
Nathalie Le Clef	ESHRE Central Office, Grimbergen, Belgium
Kersti Lundin	Sahlgrenska University Hospital, Sweden
Cristina Magli ^a	SISMER, Bologna, Italy
David Mortimer ^a	Oozoa Biomedical, Vancouver, Canada
Sharon Mortimer	Oozoa Biomedical, Vancouver, Canada
Zsolt Peter Nagy	Reproductive Biology Associates, Atlanta, USA
Johan Smitz ^a	Centre for Reproductive Medicine, Brussels, Belgium
Arne Sunde	St Olav's University Hospital, Trondheim, Norway
Nathalie Vermeulen	ESHRE Central Office, Grimbergen, Belgium

^a Presenter at the consensus workshop.

especially in ART, there is a general lack of evidence to support their importance, scientific soundness and usefulness (Shahangian and Snyder, 2009).

During the consensus meeting, the results of the surveys, scientific evidence and personal clinical experience were integrated into presentations by experts on specific topics. For each indicator, information was presented in a fixed format: definition, rationale, qualifiers, formula, data sources, KPI strengths and weaknesses, frequency of data collection and reference values for minimum expected and target values based on 50 and 75 percentile values, respectively. After the presentation for the topic, each proposed indicator was discussed until consensus was reached within the group.

After the meeting, a report was prepared describing the presentations (workshop report) and the consensus points. After approval of the report by the meeting participants, the national and international societies that contributed to the questionnaires were invited to review the report and submit comments. The final version of the manuscript was approved by the Executive Committees of ESHRE and Alpha before publication.

This paper is divided into two parts: the workshop report, and the recommendations of the Expert Panel.

Workshop report

Effects of ovarian stimulation on embryology parameters

The methods of ovarian stimulation have been evolving since the earliest days of clinical IVF, in the search for the best stimulation protocol. With that goal in mind, there has been an enormous effort to develop

the best pharmaceuticals and protocols, but in practice economic factors as well as prevailing opinion can influence treatment decisions beyond consideration of the patient's endocrine background.

Despite the thousands of smaller studies on this subject in the literature, there are very few large multicentre randomized controlled trials (RCT) beyond those organized by pharmaceutical companies, and these do not consider 'non-standard' groups of patients. Furthermore, the meta-analyses that have been conducted are often unclear about their inclusion criteria. As a result, the prevailing approach is to use what works within each clinic, and is often uncorrected for the patients' weight, body mass index (BMI), or endocrine background. Some clinics also do not monitor cycles for financial reasons, which can have an influence on the cycle outcome.

In general, a 'good' stimulation is one that produces a homogeneous cohort of mature oocytes, with the least inconvenience and risk to the patient, and results in the birth of a healthy singleton. From the Clinical Embryologist's perspective, a good stimulation results in the retrieval of well-expanded cumulus-oocyte complexes (COC), as is expected from each follicle >14 mm in diameter, with a high proportion of metaphase II (MII) oocytes (Ectors et al., 1997; Nogueira et al., 2006; Scott et al., 1989). On the other hand, a poor stimulation, caused, for example, by sub-optimal decisions regarding timing or stimulation dose, is one that may result in a high rate of abnormal COC morphology observed at the time of oocyte retrieval, possibly resulting in an increased rate of abnormal fertilization (e.g. 1 pronucleus [PN], 3PN, etc.) and/or decreased rate of normal fertilization and an increased aneuploidy rate (Soares et al., 2003).

Aggressive ovarian stimulation has effects on the patient's well-being, by increasing the risk of ovarian hyperstimulation syndrome (OHSS) (Delvigne, 2009; Rizk, 2009), as well as on the endometrium and the ovaries. There are studies showing an increased likelihood of embryo aneuploidy in connection with aggressive ovarian stimulation, even in patients younger than 35, including post-zygotic segregation errors and maternal segregation errors (Baart et al., 2007; Haaf et al., 2009), as well as having a negative impact on the maintenance of genomic imprints during early embryogenesis (Denomme and Mann, 2012; Fauque et al., 2007; Saenz-de-Juano et al., 2016). It has been shown that in patients belonging to either high or low strata of antimüllerian hormone (AMH) concentration an inverse correlation exists between the daily dose of recombinant human follicle stimulating hormone (rhFSH) used in the stimulation and the proportion of blastocyst formation (Arce et al., 2014).

To determine whether there is a stimulation method that could yield a higher number of competent oocytes, one first needs to consider the effects of LH and FSH as the principal drivers of ovarian stimulation, and their pharmacodynamics. In a natural cycle, FSH receptor expression peaks during the early follicular phase then declines, while LH receptor expression increases from mid-follicular phase, indicating that LH is likely to be involved in follicular development (Jeppesen et al., 2012). This could explain, at least in part, why follicular recruitment is compromised in women with profound pituitary downregulation (Ferraretti et al., 2004). The role of the theca cells in ovarian responsiveness to FSH is also illustrated by compromised follicular recruitment in women older than 35 years (Hugues et al., 2010; Humaidan et al., 2004; Piltonen et al., 2003), particularly those with reduced ovarian sensitivity to FSH (Davison et al., 2005) and reduced ovarian capacity to secrete androgens under basal LH conditions (Spencer et al., 2007). In addition, it has been shown that LH induces epidermal growth factor-like factors in the mural

granulosa. Amphiregulin, one of these factors, has been correlated with good oocyte quality (Sugimura et al., 2015; Zamah et al., 2010).

The time of oocyte retrieval relative to the time of the ovulation trigger is typically in the range of 34–38 h. In a meta-analysis including 5 RCT with a total of 895 women, it was found that a time of oocyte retrieval relative to the time of the ovulation trigger of >36 hours compared to <36 hours resulted in a higher oocyte maturation rate, but no difference in fertilization rate, implantation rate or pregnancy rate (Wang et al., 2011). Deviations from the locally established protocol should be documented and taken into consideration.

In conclusion, to optimize outcomes, there is a need for individualization of the treatment protocol. As stimulation can affect a range of systems, closely monitoring the patient's response could reduce many of the risks associated with ovarian stimulation, as stimulation can affect a range of systems, which will also have an impact on the success of treatment.

Because of the interlinked effects of ovarian stimulation on oocyte quality and developmental competence, it is logical that the most successful clinics are those in which the embryologists and clinicians speak with each other and communicate regularly and effectively about outcomes related to stimulation (and other procedures) (Van Voorhis et al., 2010).

Oocytes

Not all oocytes collected from a patient following ovarian stimulation for ART will have the same developmental competence, which is illustrated by the observation that only 5% of oocytes collected eventually result in a live birth (Lemmen et al., 2016). Intrinsic oocyte competence is derived not only from the degree of nuclear maturity of the oocytes, but also from their cytoplasmic maturity (Garrido et al., 2011; Lemmen et al., 2016; Patrizio and Sakkas, 2009). Furthermore, oocyte developmental competence is affected by a range of intrinsic patient-related and external factors. These patient-related factors include age, BMI, lifestyle factors, and type of infertility. The external factors include ovarian stimulation, laboratory procedures (such as oocyte retrieval, denudation, cryopreservation, preparation for ICSI), culture conditions (temperature, pH, pO₂), environmental conditions (light, air quality, humidity) and culture medium.

Against that background, the question is whether any laboratory indicator can provide a measure of the intrinsic oocyte competence at the time of oocyte retrieval, as all of the subsequent events could be influenced by laboratory procedures, and/or by the genetic contribution of the spermatozoon. In other words, is quality measurable for oocytes, or perhaps more pertinently, is there any measure that could pinpoint where a dysfunction occurred during the long process of oocyte development?

There are a number of possible markers for oocyte competence, but these are largely research-based, and have not found widespread application in clinical service. These include assessment of biochemical markers in follicular fluid, gene expression studies of follicular cells, and oxygen uptake assessments (Nagy et al., 2009; Nel-Themaat and Nagy, 2011). Other markers, such as assessment of oocyte morphology, spindle imaging, and polar body biopsy can be incorporated into clinical service, but this is not a universal approach (Braga et al., 2013; Patrizio et al., 2007; Rienzi et al., 2011).

When the results of the Alpha and ESHRE surveys were combined, respondents identified oocyte recovery rate and oocyte maturity rate as the most important indicators for oocytes. Although, strictly speaking, they do not provide an indication of laboratory performance,

they do provide an estimate of response to stimulation, and therefore a general estimation of the likely developmental competence of the oocytes.

In the Alpha survey, oocyte recovery rate was defined as the likelihood of aspirating a COC from each follicle over a certain size as measured on the day of triggering. The rationale for this is the expectation that those follicles that have achieved a certain size, with a good response to FSH and a sufficient number of LH receptors in follicular cells, will respond appropriately to the ovulation signalling cascade, resulting in the release of the COC into the follicular fluid, thereby facilitating its aspiration. A concern with this potential indicator was its reliance on the accuracy of follicular scanning, and the need for a consistent time interval between ovulation trigger and oocyte retrieval. However, while a range of follicle sizes were identified in the survey as the 'ideal' size for triggering, the expected recovery rates were remarkably similar, generally ranging from 70–80% as the competence level, and 85–100% as the benchmark value.

Oocyte maturity rate is generally related to nuclear maturity, being defined as the proportion of oocytes at MII stage. Its potential value is as a marker of the efficiency of ovarian stimulation and triggering. Of the Alpha survey respondents, 80% indicated that their laboratory determined the MII rate, with median competence and benchmark values of 75% and 90%, respectively. It was noted that the timing of this assessment is an important factor, as it is not possible to assess oocyte maturity at the time of oocyte retrieval in the case of insemination by routine IVF. Since this assessment requires the removal of the cumulus and corona cells, it can be performed at the time of denudation of the oocytes prior to ICSI, but for a universal competence and benchmark value to be established, a consistent time interval between the time of trigger and the time of cumulus cell removal would be required (e.g. 40 ± 1 h post ovulation triggering).

A third potential indicator, oocyte grade, was defined as the proportion of COC with expanded cumulus at the time of oocyte retrieval. As ovulation triggers cumulus expansion by mediating the synthesis of hyaluronic acid and the organization of a stable cellular matrix [Russell and Salustri, 2006], this indicator provides an assessment of the quality of communication between the oocyte and its cumulus cells. The potential problems associated with the use of this indicator are a lack of objective criteria for making this assessment, and a concern that there is not always a good correlation between nuclear maturity and cumulus cell expansion [Balaban and Urman, 2006].

Other indicators that were proposed, but not considered valuable or reliable by the survey respondents were: rate of degenerated (or empty) zona pellucidae; rate of germinal vesicle (GV) oocyte recovery; definition of the minimum number of follicles to justify flushing; and oocyte degeneration rate at the time of oocyte retrieval. Oocyte degeneration rate after removal of cumulus cells is discussed in the section on ICSI.

Overall, the responses to the surveys highlighted the lack of consistent data for the evaluation of oocyte quality and competence, and identified an opportunity for national and international registries to promote the collection of this information.

Spermatozoa

Proposed andrology laboratory PIs were sperm recovery rate, and sperm motility post-wash. In addition, sperm parameters were discussed in relation to the decision for intrauterine insemination (IUI), IVF or intracytoplasmic sperm injection (ICSI).

Survey responses revealed such wide ranges in perceived semen analysis minima for suitability for IUI, IVF or ICSI, as well as expected sperm recovery post-wash, as to make it impossible to determine robust recommendations for competency and benchmark values for any of these criteria.

With regards to sperm preparation, it is possible that respondents were confused when reporting the 'recovery rate', and the substantial variability in terms of the expected/required number of spermatozoa in the final preparation likely included confusion between % recovery ('yield') values and the actual number of spermatozoa (millions); hence these data were considered unreliable. However, the expected proportion of motile spermatozoa in the final washed preparation showed coherence across the respondents, with both median and mode values of 90% for competency and 95% for the benchmark.

A major issue when considering semen analysis data is that many ART laboratories do not employ methods that meet the minimum standards required by either the World Health Organization (WHO) or the ESHRE SIG Andrology [Björndahl et al., 2010; World Health Organization, 2010]. Therefore, reported values for sperm concentration and motility must be understood to have high uncertainty of measurement, and hence need to be considered as inaccurate and unreliable [Björndahl et al., 2016; Sanchez-Pozo et al., 2013]. As a consequence, any association between semen analysis characteristics, yield and fertility potential will remain unclear if based on studies using inappropriate semen analysis techniques. From a best practice standpoint, any clinical laboratory providing semen analysis or post-preparation values that are to be used for diagnostic or treatment management purposes should participate in an external quality assurance (EQA) programme which provides a comparison between the participating laboratories' results and established reference ('correct answer') values so as to permit quality improvement in laboratory work [Björndahl et al., 2010].

Moreover, there is a general concern that semen analysis reference values have little or no value for ART procedures [Björndahl, 2011]. In particular, the WHO reference values for sperm concentration, motility and vitality were derived from populations of men who had achieved in-vivo conceptions [Cooper et al., 2010], and therefore these cut-off values have no *a priori* relevance in regard to ART patients, and hence the need or suitability for any form of ART treatment should not be decided based on these reference values.

The Tygerberg Strict Criteria for normal sperm morphology were derived in regard to ART success [Coetzee et al., 1998; Kruger et al., 1988], so these cut-off values might be pertinent in differentiating between the need for IUI, IVF or ICSI – although concern regarding measurement uncertainty cannot be ignored [Menkveld et al., 2011]. While a cut-off of 4% normal forms might help define sub-populations of patients with differing prognoses, at the level of individual patients a result of 4% based on 200 spermatozoa evaluated is not very informative since the result has a statistical expectation ranging from 2–8% [Björndahl et al., 2010], and to be able to differentiate between 3% and 5% with statistical robustness would require the assessment to have been made evaluating over 1500 spermatozoa.

Because of the limitations of semen analysis cut-off values, a decision on the suitability of IUI or routine IVF for a couple should be made based on post-preparation sperm number/concentration and motility, ideally assessed during a pre-treatment 'trial wash' (while still taking into account the uncertainty of measurement). In case of ICSI, there should not be any cut-off based on semen analysis characteristics, the only logical criterion would be having sufficient (in comparison with the expected number of oocytes) spermatozoa

that are, ideally, viable, and preferably motile or hypo-osmotic swelling (HOS) test positive [Nagy et al., 1995]. As a general principle, laboratories should develop and apply their own criteria for deciding on IUI, IVF or ICSI, based on the couple's clinical situation and reproductive history rather than on semen analysis.

Regarding andrology laboratory PIs, results from the Alpha survey indicated that only post-preparation sperm motility would be a valuable indicator, as it monitors the effectiveness of the sperm washing procedure. Therefore, post-wash sperm motility should be monitored for fresh ejaculate specimens that show normozoospermia as per the WHO5 guidelines [World Health Organization, 2010], but still taking into account the poor reliability of sperm motility data, non-robust classification of semen samples based on the high uncertainty of semen analysis data, and variability in sperm preparation methods.

Sperm recovery rate, defined as the percentage recovery of progressively motile sperm after washing as compared to pre-washing [Björndahl et al., 2010], can be used as a laboratory KPI, providing useful information for inter-operator comparison and proficiency testing. However, given the high uncertainty in counting and the different protocols for sperm preparation (notably with density gradient washing resulting in higher recovery rates compared with direct swim-up from semen), no competence values can be provided. Laboratories should develop their own standards according to their own clinical and laboratory practice.

Fertilization after insemination by ICSI

Although several potential KPIs have been identified in ICSI, the presentation focused on the four most pertinent: normal fertilization rate, oocyte degeneration rate, poor fertilization rate and failed fertilization rate.

The definition used most often for the ICSI normal fertilization rate is the proportion of injected oocytes with 2PN the day after injection, except for the Spanish Registry and the Istanbul Consensus which include the observation of two polar bodies (PB) in the definition [Alpha Scientists In Reproductive Medicine and ESHRE Special Interest Group of Embryology, 2011]. The suggested competence and benchmark values for this indicator were consistent among respondents ranging from 60–80% and 70–100%, respectively. The UK's Association of Clinical Embryologists (ACE) proposed benchmark for the 2PN rate is >65%, including only patients below 40 years of age with at least three oocytes collected [Hughes and Association of Clinical Embryologists, 2012]. From the literature it was found that ICSI results in an average fertilization rate of 70% [Heindryckx et al., 2005; Kashir et al., 2010], which was similar to 68.7% based on over 20,000 unselected MII oocytes at the CARE Fertility laboratories (personal communication, Alison Campbell). ICSI fertilization rate is a commonly reported and effective indicator that is informative of gamete quality and operator competence. ICSI 2PN rate does depend on the various criteria used for performing ICSI, which can be considered a weakness of the indicator.

The ICSI damage rate or oocyte degeneration rate was ranked as important. In the Alpha survey, the minimum expected value and target value ranged from 3–30% and 0–10%, respectively. Oocyte damage can be observed at three time points during the ICSI process from the start at stripping, during ICSI, or at the fertilization assessment on Day 1. Oocyte damage probably occurs most frequently during injection, but without immediate signs of damage this is not detected until the fertilization check. In addition, as both the damage detected at injection and at fertilization check reflect damage from the

ICSI process, these should not be recorded and calculated separately. Damage at denudation/stripping can be monitored separately as it mainly reflects operator competency, but it has a very low frequency. ICSI damage rate is therefore defined as the number of oocytes damaged during ICSI, and/or observed at fertilization check over the number of injected oocytes. It is useful to monitor this indicator for operator competence, oocyte quality, and laboratory performance. The damage rate can also be indicative of technical problems (e.g. cumulus cell removal stress, vibration). Alternatively, the term 'ICSI oocyte survival rate' can be used.

Poor fertilization rate is defined as the proportion of cycles in which <25% of the injected oocytes are fertilized. The responses from the survey are much divided, ranging from 5%–20% for the competence and from 0–15% for the benchmark value. Poor fertilization rate can give an indication of operator competence and reflect gamete quality.

Failed fertilization rate scored relatively low in importance in the surveys. Failed fertilization rate is defined as the proportion of cycles in which none of the injected oocytes are fertilized. The indicator can be informative of gamete quality/function and/or operator skill. A deficiency in the mechanism of oocyte activation is regarded as the principal cause of ICSI fertilization failure or abnormally low fertilization. Complete (or 'virtually complete') fertilization failure with ICSI occurs in 1–5% of cycles [Flaherty et al., 1998; Kashir et al., 2010; Liu et al., 1995; Mahutte and Arici, 2003; Yanagida, 2004]. From the Alpha survey there was a wide range in the competence values for this indicator ranging from 2–15% (median 5%), with a benchmark of <1%, respectively.

For these and other KPIs, a reference population could be relevant. With regard to ICSI fertilization rate, it could be relevant to exclude cases where reduced fertilization rates are anticipated, including in-vitro matured MI oocytes (although inconclusive data), artificially activated oocytes, use of testicular sperm, and cases of globozoospermia and asthenozoospermia [Rubino et al., 2016]. However, due to the low prevalence, including these cases may not significantly affect the indicators in most clinics.

In conclusion, from the surveys and collected evidence, ICSI damage rate and ICSI normal fertilization rate are considered relevant and important KPIs, while the value of ICSI low/failed fertilization rate as a KPI is less clear. Oocyte maturity rate and 1PN/3PN rate were not specifically discussed for ICSI. The ICSI rate, defined as the proportion of cycles that use ICSI, was not commonly recorded by the survey respondents, judging this as a less relevant PI.

Fertilization after (routine) IVF insemination

This section deals with normal fertilization rate, polyspermy rate, poor fertilization rate, and zygote morphology after routine IVF insemination.

Pronuclear formation occurs 1.5–2.0 hours earlier in oocytes inseminated by ICSI compared with those inseminated by conventional IVF [Montag et al., 2001; Nagy et al., 1998]. This should be taken into consideration when setting the time for fertilization check, relative to the time elapsed since insemination (recommended as 17 ± 1 h) [Alpha Scientists In Reproductive Medicine and ESHRE Special Interest Group of Embryology, 2011]. A normally fertilized oocyte should have two pronuclei (2PN) of similar size that are closely apposed and centrally located [Alpha Scientists In Reproductive Medicine and ESHRE Special Interest Group of Embryology, 2011]. Although the provenance of micronuclei remains unclear, their presence could be considered to be abnormal, as could the presence of pronuclei of different sizes. From the literature, the normal fertilization rate (i.e. the

proportion of inseminated oocytes with 2PN at the time of the fertilization check on Day 1) is expected to be around 67%, with a range of reported values from 53–81%. This range was reflected in the Alpha survey results, where the median value for competency was 61% of inseminated oocytes, with a median benchmark value of 77%.

The presence of three or more pronuclei is indicative of an abnormal fertilization, arising from either failure to extrude the second polar body or polyspermy. Polyspermy may be the result of either oocyte immaturity (causing failure of the cortical reaction), oocyte overmaturity, and/or an extremely high concentration of motile spermatozoa in the insemination volume (Wang et al., 2003). From the literature, the incidence of ≥ 3 PN is 4–7% in IVF (Joergensen et al., 2015). This agrees well with the median values from the Alpha survey of <9% for competence, and <4.5% as a benchmark.

The calculation of the total fertilization rate following IVF includes all fertilized oocytes with ≥ 2 PN. Although, as already stated, oocytes with >2 PN are abnormally fertilized, this parameter provides an indication of the ability of the culture system to support sperm capacitation and sperm–oocyte interaction in IVF cycles. Of the published studies that included >100 oocytes, the median total fertilization rate was 76% (range 69–87%) in selected patient populations. This is similar to the Alpha survey results which suggested a competency level of at least 63% and a benchmark of at least 84%.

Oocytes with a single pronucleus after insemination by IVF, which occurs in 1–5% of cases, can be indicative of fertilization and syngamy, asynchronous appearance of pronuclei (an extremely rare event, as evidenced by the use of time-lapse microscopy), or parthenogenetic activation (Levron et al., 1995). The incidence of diploidy in 1PN oocytes following conventional IVF has been reported to be in the range of 45–50% (Kai et al., 2015; Staessen and Van Steirteghem, 1997; Sultan et al., 1995). In contrast, 1PN oocytes arising after ICSI have a reported diploidy rate of only 7–14%, with genetic abnormalities in the subsequent embryos (Mateo et al., 2013).

The incidence of poor fertilization (<25% of inseminated COC with 2PN) or total failure of fertilization (no oocytes with signs of fertilization) could be indicative of a problem with sperm function, too few motile spermatozoa during insemination, or failure of oocyte activation (Ebner et al., 2015). There is very little evidence in the literature regarding the expected incidence of either poor or failed fertilization. However, the Alpha survey results suggested competency and benchmark levels for poor and failed fertilization of 14% and 6%, and 8% and 4%, respectively.

Although the Istanbul Consensus made recommendations about grading zygote morphology (Alpha Scientists In Reproductive Medicine and ESHRE Special Interest Group of Embryology, 2011), the question remains as to whether indicators referring to zygote morphology are useful, especially as differences in pronuclear pattern could be related to the insemination method and timing of the observation (Ebner et al., 2003; Montag et al., 2001).

Cleavage-stage embryos

Proposed indicators for cleavage-stage embryos are early cleavage rate, cleavage rate, embryo development rates, embryo fragmentation rate, and rate of good quality embryos (embryo score or grade).

Early cleavage rate is defined as the proportion of cleaved zygotes at the early cleavage check on Day 1 (26 ± 1 h post-ICSI or 28 ± 1 h post-IVF) (Alpha Scientists In Reproductive Medicine and ESHRE Special Interest Group of Embryology, 2011), but other time intervals after insemination have been used for assessing early cleavage.

This indicator reflects the ability of the culture system to support early cleavage of fertilized oocytes and the viability and quality of the embryos (Shoukir et al., 1997). There are conflicting results on the importance of early cleavage. Studies have shown that early cleavage, together with other factors, can be used as an embryo selection method (Ciray et al., 2005; Lundqvist et al., 2001). Early cleavage rate has also been shown to correlate with blastocyst implantation and pregnancy rates (Balaban and Urman, 2003; Shoukir et al., 1997) and it is a better independent marker of implantation potential than zygote morphology (Brezinova et al., 2009). In contrast, early cleavage was not found to be an independent predictor of implantation in IVF patients with good prognosis (Thurin et al., 2005). In addition, premature occurrence of early cleavage can be negatively, instead of positively, associated with embryo implantation potential (Meseguer et al., 2011). Furthermore, it was not a reliable predictor for embryo implantation rate when good quality embryos are transferred (de los Santos et al., 2014; Sundstrom and Saldeen, 2008), or when using a GnRH antagonist protocol (Yang et al., 2015). In the Alpha survey, competence and benchmark values ranged from 10–50% and 15–60%, respectively.

Cleavage rate reflects the ability of the culture system to support cellular division of fertilized oocytes. It is an indicator of embryo viability and has the ability to detect culture media contaminants. The presence of non-cleaved embryos or arrested zygotes on Day 3 is associated with a decrease in quality of the remaining cohort, but without a negative impact on clinical outcome (Machtinger et al., 2015). Cleavage rate is considered important and widely monitored, and defined as the proportion of zygotes which cleave to become embryos on Day 2 at 44 ± 1 h post-insemination (Alpha Scientists In Reproductive Medicine and ESHRE Special Interest Group of Embryology, 2011). Cleavage rate should be calculated not only in the total population, but also in reference groups (IVF versus ICSI, female age, ejaculated versus surgically retrieved sperm), and controlled for confounders (the timing of observation and oocyte maturity). Also, the presence of a refractile body in the oocyte is associated with reduced cleavage rates and impaired embryo development, while the cytoplasmic granularity did not seem to have an effect on embryo development (Fancovits et al., 2012). Cleavage rate should be calculated frequently (at least once per month). In the Alpha survey the competence values ranged from 80–95%, with a benchmark of 90–100%.

Embryo development rate is defined as the proportion of 4-cell embryos on Day 2 among the 2PN zygotes (measured at 44 ± 1 h post-insemination), the proportion of 8-cell embryos on Day 3 (measured at 68 ± 1 h post-insemination), and the proportion of morula-stage embryos on Day 4 (92 ± 2 h post-insemination) (Alpha Scientists In Reproductive Medicine and ESHRE Special Interest Group of Embryology, 2011). This indicator reflects the ability of the culture system to support cleavage according to the expected developmental stages and the quality and viability of embryos, especially for Day 2 or Day 3 transfer, while less important for blastocyst transfer. Possible confounders are the timing of laboratory observations and the type of culture media used. Although dependent on iatrogenic factors like the culture conditions, embryo development rate is an important indicator; in well-defined categories of patients, it reflects the overall laboratory performance. It was stated that sufficient numbers of embryos or longer data collection are necessary, as this indicator is influenced by short-term variations. Calculation of a sliding mean can be helpful for detecting long-term variations. The value and practicability of the morula check on Day 4 was questioned, as centres performing Day 5 embryo transfer often do not assess the embryo

development at Day 4. Also, instead of assessing the number of 4-cell embryos on Day 2, or 8-cell embryos on Day 3, a combination of 4-cell and 8-cell embryos on Day 2 and Day 3 can be used. This was analysed in a study by van Royen, who thereby characterized a top-quality embryo as having 4–5 cells on Day 2 and ≥ 7 cells on Day 3 (Van Royen et al., 1999). In larger centres, assessing the embryos at the specified time points can be difficult, and counting 4- and 5-cell embryos on Day 2, and 7-, 8- and 9-cell embryos on Day 3 may be more relevant.

The rate of good-quality embryos is defined as the proportion of Day 2 and Day 3 embryos with high score or grade. Many different scoring systems exist, based on different variables, including cell number, fragmentation, cell size and multinucleation (Alpha Scientists In Reproductive Medicine and ESHRE Special Interest Group of Embryology, 2011) and blastomere nuclear status (Fauque et al., 2013). A recent study evaluating which set of embryo variables is most predictive for live birth rate (LBR) reported that blastomere number, proportion of mononucleated blastomeres, degree of fragmentation, and variation in blastomere size were significantly associated with LBR in univariate analysis, while symmetry of the embryo was not (Rhenman et al., 2015). Furthermore, the grading systems are not robust, but can be used as internal quality assessment parameters. The importance of this parameter is also affected by the limited significance of the fragmentation rate.

Embryo fragmentation rate, defined as the proportion of Day 2 and Day 3 embryos with $<10\%$ fragmentation, reflects the quality and viability of embryos. From the Alpha survey the competence value for this parameter ranged from 20–90% (median 50%) and the benchmark from 30–90% (median 70%). These large ranges underline the difficulties with this parameter: embryo fragmentation rate is reported to be a subjective parameter and difficult to evaluate as one has to differentiate between a cell and a fragment and then estimate the relative proportion of fragments (Paternot et al., 2011).

Embryo utilization rate is defined as the number of embryos utilized (transferred or cryopreserved) per number of 2PN zygotes in the same cycle. This parameter is often presented in studies, but competence and benchmark values cannot be calculated due to its dependence on strategies for embryo transfer and cryopreservation, as well as patient request.

In conclusion, embryo cleavage rate and embryo development rate are extremely important indicators, while early cleavage rate, rate of good-quality embryos, and embryo fragmentation rate are less important as quality indicators.

Blastocyst development

In the case of blastocyst-stage embryo transfer, several parameters were suggested as indicators: blastocyst development rate, good blastocyst development rate, the proportion of good quality blastocysts, and Day 5 embryo transfer rate.

The blastocyst development rate, defined as the proportion of 2PN zygotes (not just of cleaved zygotes) which are at the blastocyst stage at Day 5 (116 ± 2 h post-insemination) (Alpha Scientists In Reproductive Medicine and ESHRE Special Interest Group of Embryology, 2011), was rated important because it reflects the efficiency of the whole culture system. Blastocyst development rates can be calculated on Day 5, Day 6 or Day 5/Day 6 combined. Assessment and calculation on Day 5, consistent with previous consensus, is preferred based on limited numbers of embryos available on Day 6. The competence and benchmark values for blastocyst development rate on Day 5 ranged from 25–60% and 44–80%, respectively. The blastocyst development rate

is an objective parameter, but dependent on the assessment of blastocyst morphology, which is straightforward in the case of good quality blastocysts, but can be challenging for embryos showing an attempted cavitation. Confounders can be the timing of laboratory observation, the culture medium and the culture conditions (in particular the pO_2).

The good blastocyst development rate is defined as the proportion of 2PN zygotes which are good-quality blastocysts on Day 5 (116 ± 2 h post-insemination) (Alpha Scientists In Reproductive Medicine and ESHRE Special Interest Group of Embryology, 2011). Similar to blastocyst development rate, measuring this as an indicator only on Day 5 could make the indicator more robust. Blastocyst quality should be based on (i) blastocoele expansion, (ii) appearance of trophoctoderm (TE), and (iii) appearance of inner cell mass (ICM). Although all three parameters have been shown to be correlated to pregnancy and LBR (Van den Abbeel et al., 2013), only TE was found to be a statistically significant independent predictor of live birth outcome after adjustment for known confounders (Ahlgren et al., 2011). Even though ICM is important, a strong TE layer is essential at this stage of embryo development, allowing successful hatching and implantation (Ahlgren et al., 2011). In the Alpha survey, the competence and benchmark values for good blastocyst development rate on Day 5 ranged from 15–45% and 25–80%, respectively.

The proportion of good-quality blastocysts can be calculated from the blastocyst development rates and the good-quality blastocyst development rates. There is no evidence pertaining to the significance of this parameter in the literature, and no data from the Alpha survey.

Day 5 embryo transfer (ET) rate was defined as the proportion of cycles with ≥ 1 2PN zygotes on Day 1 that had ≥ 1 blastocyst for transfer on Day 5. From the Alpha survey there was a large variation in the values for competence and benchmarks (ranging from 25–90% and 40–100%, respectively), which is assumed to be due to differences in the time of blastocyst assessment, and different grade of expansion. In addition, this parameter depends on different policies for transfer in different settings. Cycles for preimplantation genetic diagnosis (PGD) or preimplantation genetic screening (PGS) – globally indicated as preimplantation genetic testing (PGT), comprising PGD for single gene disorders or for chromosome structural abnormalities, and PGS for aneuploidy – should be excluded from this calculation. Some possible confounders are the timing of laboratory observation (ideally at 116 ± 2 h post-insemination), but in some laboratories the time of observation depends on the timing of ET (physician availability), the culture medium and other culture conditions (e.g. pO_2), and the grade of blastocyst expansion.

Fresh cleavage or fresh blastocyst transfer?

Recently, a Cochrane review summarized evidence from RCT comparing the reproductive outcomes after fresh cleavage-stage versus fresh blastocyst-stage embryo transfer (Glujovsky et al., 2016). Based on low-quality evidence, they concluded that the LBR following fresh transfer was higher in the blastocyst-stage transfer group compared with the cleavage-stage transfer group (OR 1.48, 95% CI 1.20–1.82; 13 RCT, 1630 women, $I^2 = 45\%$). This translated to a LBR of 29% after fresh cleavage-stage transfer, and between 32% and 42% after fresh blastocyst-stage transfer. The 5 RCT that reported cumulative pregnancy rates after fresh and frozen transfers showed no significant difference after one oocyte retrieval (based on very low-quality evidence). The main limitation was serious risk of bias, associated with failure to describe acceptable methods of randomization, and unclear or high risk of attrition bias (Glujovsky et al., 2016).

Another review reported that blastocyst (Day 5/6) transfer in a fresh IVF/ICSI treatment cycle significantly increased LBR (OR 1.77; 95% CI 1.32–2.37), clinical pregnancy rate, implantation rate, and ongoing pregnancy rate, and reduced first trimester miscarriage rate, in comparison with cleavage-stage (Day 2/3) embryo transfer based on 7 RCT and 1446 cases (Wang and Sun, 2014).

Implantation rate and live birth rate

Implantation rate is judged an important indicator that reflects the overall performance of the laboratory, and an overall low implantation rate is a serious alert. Implantation rate is defined as the number of gestational sacs observed divided by the number of embryos (cleavage-stage or blastocysts) transferred (Zegers-Hochschild et al, 2009), or as the proportion of fetal heartbeats detected relative to the number of embryos transferred (Alpha Scientists In Reproductive Medicine, 2012). Implantation rate reflects the efficiency of the whole culture system, but it can be influenced by uterine receptivity, and by different policies for embryo transfer in different centres.

LBR may be considered as the ultimate KPI for checking IVF clinic performance and is defined as the likelihood of a baby being born per embryo transferred. LBR is largely affected by a series of clinical maternal factors pertaining to post-implantation development, rather than reflecting laboratory performance. This parameter can be calculated annually, but it is often difficult to collect the data.

Preimplantation genetic testing (preimplantation genetic diagnosis/screening)

The indicators proposed for PGD/PGS (PGT) were successful biopsy rate/tubing rate, rate of no biopsy, proportion of samples submitted to analysis where no results were available (no DNA was detected), embryos not found on warming. Other indicators suggested in the survey were 'survival after warming', embryo transfer per PGT cycle, and miscarriage rate, but there were very few responses for these variables in the Alpha survey.

The successful biopsy rate is defined as the proportion of biopsied and tubed/fixed samples where DNA is detected. It was suggested that this combined parameter be used – combining the tubing rate (the proportion of embryos where cells were tubed) and the proportion of samples submitted to analysis where no DNA was detected – as the tubing of cells can often not be inspected visually and will only be detected by the presence of DNA after amplification. A benchmark for the tubing rate of 95% was reported in the Alpha survey. Confounders for the successful biopsy rate are embryo quality, and the criteria for biopsy (for Day 3 embryos: presence of a visible nucleus; for Day 5/6: grade of hatching, TE quality). From the data of the ESHRE PGD Consortium, of 254,820 samples that were biopsied, 91.3% were diagnosed (De Rycke et al., 2015). It is important to underline that, as already mentioned, these data mainly refer to Day 3 biopsies and that, in the case of PGT for aneuploidy (generally known as PGS) the analysis of biopsies was mainly done by Fluorescence In Situ Hybridization (FISH), a technique requiring cells to be fixed on a glass slide. FISH is now being replaced by other methods providing 24-chromosome testing, which require tubing of cell biopsies.

The rate of no biopsy is defined as the proportion of intended PGD/PGS (PGT) cases where there were no embryos available to biopsy. This parameter was rated important, but it reflects patient-related factors and the ability of the culture system to support cleavage/blastocyst formation, rather than the performance ability of the

laboratory to perform a PGD/PGS (PGT) treatment/analysis. Furthermore, the parameter is different whether biopsy is performed at the cleavage or blastocyst stage, and depends on the timing of laboratory observation, culture medium and culture conditions (e.g. pO₂), criteria for biopsy, the time point of assisted hatching (Day 3 or Day 5), and patient selection. The parameter should also be calculated for PGD and PGS separately, based on the difference in patient populations. From the Alpha survey, the median competence value and benchmark were 20% and 10%, respectively. In the last data collection of the ESHRE PGD Consortium, out of 45,163 reported cycles, 2.8% were cancelled before biopsy (De Rycke et al., 2015). However, as this data collection covers cycles performed up to 2010, this figure refers to Day 3 biopsies, implying that possibly a higher value could be expected for Day 5 biopsies.

For a number of samples, no results are available after amplification. Results can be inconclusive even with a strong DNA amplification band in the agarose gel, which can be due to early fragmentation of DNA. In single gene analysis, results can be inconclusive if information is found only on a reduced number of markers, or the gene of interest failed to amplify. Although this indicator is not strictly related to the performance of the laboratory, it is relevant to inform clinicians of the total number of embryos where PGD/PGS (PGT) results are available. The number of samples where no results are available after amplification was reported in the Alpha survey with a benchmark of <5%, which corresponds with the converse of the tubing rate.

Nowadays, the majority of PGT cycles are based on blastocyst biopsy and cryopreservation. The proportion of embryos not found on warming and embryos degenerated after warming mainly reflect operator skill. In the Alpha survey, the median competence values were 3% and 10% for not-found and degenerated embryos, respectively. These indicators can also be reported as one combined indicator. It was mentioned that clinicians should be aware that although laboratories strive for 100% recovery, not all embryos submitted to PGD/PGS (PGT) will be recovered after warming.

Reference values for implantation rate and LBR after PGT can be derived from the data from the ESHRE PGD Consortium. The implantation rate was 26% for PGD and 22% in PGS, with delivery rates (per oocyte retrieval) of 21% and 14%, respectively (De Rycke et al., 2015).

Indicators for cryopreservation: addition to the previous consensus (Alpha Scientists In Reproductive Medicine, 2012)

Blastocyst re-expansion rate is defined as the proportion of warmed blastocysts that show re-expansion within a defined time period (e.g. 2 h). Recent evidence shows that there is an impact on the performance results depending on the quality/expansion of the blastocysts which are cryopreserved (Cobo et al., 2012). Also, in blastocyst fresh transfer, multivariate analysis showed that the odds of live birth increased by 36% for each grade of expansion ($P = 0.0061$) and decreased by 29% for blastocysts with grade B TE compared with grade A TE ($P = 0.0099$). Furthermore, after thawing, the odds of live birth increased by 39% ($P = 0.0042$) for each 10% increase in degree of re-expansion. Therefore, blastocoel expansion and TE grade were selected as the most significant pre-freeze morphological predictors of live birth and degree of re-expansion was selected as the best post-thaw parameter for prediction of live birth (Ahlstrom et al., 2013). Confounding factors are time of observation, female age and fertilization

method. These observations do not include embryos that had been biopsied on Day 3 as they have a different hatching dynamic (Lopes et al., 2015).

Recommendations of the expert panel

General comments

- Regarding frequency of data collection for indicators, it was the consensus opinion that this should be done, ideally, on a monthly basis. However, it was recognized that this is not always practical, based on caseload, and therefore either a longer timeframe or a specific predetermined number of cases might be used instead. The minimum number will depend on the stability of the indicator and will need to be developed by the laboratory, although an initial dataset of 30 cases could be used as a guideline. Nonetheless laboratories should remain vigilant and respond promptly to unexpected fluctuations.
- The discussions identified three different types of indicator: reference indicators (RIs), PIs, and KPIs:

RIs were related to the oocytes coming into the laboratory, and so were proxy indicators of the response to ovarian stimulation. PIs were those for which data should be documented and stored, even if they are not routinely reported in a control chart. KPIs were those related to the 'core business' of the ART laboratory.
- The values for indicators are presented as competency and benchmark values, as was done for the cryopreservation consensus (Alpha Scientists In Reproductive Medicine, 2012). The gap between the competency and the benchmark values is the 'desirable range'.
- It was the opinion of the Expert Panel that ovarian stimulation can have an impact on the overall treatment cycle but is less likely to have an impact on any single laboratory PI.
- To apply the recommended values:

The time of oocyte retrieval relative to the time of the ovulation trigger is typically in the range of 34–38 h (most commonly 36h). Deviations from the locally established protocol should be documented and taken into consideration.

Timing of all observations should be made as recommended in the Istanbul consensus (Alpha Scientists In Reproductive Medicine and ESHRE Special Interest Group of Embryology, 2011)
- Individual clinics should decide whether it is more practical to subdivide their results into specific patient groups for KPI and PI

Table 2 – Reference indicators.

Reference Indicator	Calculation	Benchmark Value
Proportion of oocytes recovered (stimulated cycles)	$\frac{\text{no. oocytes retrieved}}{\text{no. follicles on day of trigger}} \times 100$	80–95% of follicles measured
Proportion of MII oocytes at ICSI	$\frac{\text{no. MII oocytes at ICSI}}{\text{no. COC retrieved}} \times 100$	75–90%

COC = cumulus-oocyte complexes; MII = metaphase II.

determinations, based on their clinical practice. The Indicator values presented here were derived relative to cycles that met the criteria for a 'reference population'. With the exception of Indicators with specific qualifiers identified, these criteria were:

female patients <40 years old;

own fresh oocytes;

ejaculated spermatozoa (fresh or frozen);

no PGD/PGS (PGT); and

all insemination methods (i.e. routine IVF and ICSI).

- It was the opinion of the panel that national and international registries should be encouraged to gather data that can be used for the derivation of KPI standard values.
- Any selection procedure in addition to embryo morphology, e.g. PGD/PGS (PGT) or time-lapse imaging, is not expected to increase the cumulative pregnancy/LBR, but in the case of PGS it may result in reduced time to pregnancy/live birth.

Indicators

The results of the discussions have been summarized for the majority of the Indicators. The values for Indicators have also been presented in Tables 2–4, but these should be read in association with the summary for each Indicator.

Proportion of oocytes recovered

This RI (Table 2) is defined as the number of oocytes retrieved as a function of the number of ovarian follicles seen at ultrasound assessment. It is useful as a measure of whether the quantity of oocytes is maximized. The values are not influenced by laboratory practice, and so cannot be held to be laboratory PI, but values outside the expected range could prompt an investigation of any changes in stimulation practice. Having this information is an important factor

Table 3 – Performance indicators.

Performance Indicator	Calculation	Competency value	Benchmark value
Sperm motility post-preparation (for IVF and IUI)	$\frac{\text{progressively motile sperm}}{\text{all sperm counted}} \times 100$	90%	≥95%
IVF polyspermy rate	$\frac{\text{no. fertilized oocytes with } > 2\text{PN}}{\text{no. COC inseminated}} \times 100$		<6%
1PN rate (IVF)	$\frac{\text{no. 1PN oocytes}}{\text{no. COC inseminated}} \times 100$		<5%
1PN rate (ICSI)	$\frac{\text{no. 1PN oocytes}}{\text{no. MII oocytes injected}} \times 100$		<3%
Good blastocyst development rate	$\frac{\text{no. good quality blastocysts on Day 5}}{\text{no. 2PN/2PB oocytes on Day 1}} \times 100$	≥30%	≥40%

COC = cumulus-oocyte complexes; ICSI = intracytoplasmic sperm injection; IUI = intrauterine insemination; PB = polar body; PN = pronucleus.

Table 4 – Key performance indicators.

Key performance indicator	Calculation	Competency value	Benchmark value
ICSI damage rate	$\frac{\text{no. damaged or degenerated}}{\text{all oocytes injected}} \times 100$	≤10%	≤5%
ICSI normal fertilization rate	$\frac{\text{no. oocytes with 2PN and 2PB}}{\text{no. MII oocytes injected}} \times 100$	≥65%	≥80%
IVF normal fertilization rate	$\frac{\text{no. oocytes with 2PN and 2PB}}{\text{no. COC inseminated}} \times 100$	≥60%	≥75%
Failed fertilization rate (IVF)	$\frac{\text{no. cycles with no evidence of fert'n}}{\text{no. of stimulated IVF cycles}} \times 100$		<5%
Cleavage rate	$\frac{\text{no. cleaved embryos on Day 2}}{\text{no. 2PN/2PB oocytes on Day 1}} \times 100$	≥95%	≥99%
Day 2 embryo development rate	$\frac{\text{no. 4-cell embryos on Day 2}}{\text{no. normally fertilized oocytes}^a} \times 100$	≥50%	≥80%
Day 3 embryo development rate	$\frac{\text{no. 8-cell embryos on Day 3}}{\text{no. normally fertilized oocytes}^a} \times 100$	≥45%	≥70%
Blastocyst development rate	$\frac{\text{no. blastocysts Day 5}}{\text{no. normally fertilized oocytes}^a} \times 100$	≥40%	≥60%
Successful biopsy rate	$\frac{\text{no. biopsies with DNA detected}}{\text{no. biopsies performed}} \times 100$	≥90%	≥95%
Blastocyst cryosurvival rate	$\frac{\text{no. blastocysts appearing intact}}{\text{no. blastocysts warmed}} \times 100$	≥90%	≥99%
Implantation rate (cleavage stage) ^b	$\frac{\text{no. sacs seen on ultrasound}^c}{\text{no. embryos transferred}} \times 100$	≥25%	≥35%
Implantation rate (blastocyst stage) ^b	$\frac{\text{no. sacs seen on ultrasound}^c}{\text{no. blastocysts transferred}} \times 100$	≥35%	≥60%

ICSI = intracytoplasmic sperm injection; MII = metaphase II; PB = polar body; PN = pronucleus.
^a Defined as oocytes with 2PN and 2PB on Day 1.
^b Based on total number of embryos transferred to all patients in the reference group, not just to those for whom an implantation occurred.
^c Definition reached after discussion, as some felt that no. fetal heartbeat detected/no. embryos transferred was a more meaningful Indicator.

in troubleshooting. The expected range is 80–95% of follicles measured in stimulated cycles.

Proportion of MII oocytes at ICSI

This RI (Table 2) is defined as the proportion of oocytes that have nuclear maturity at the time of injection, and so acts as a proxy indication of the effectiveness of ovarian stimulation. It is not a laboratory PI, as values are not influenced by laboratory practice, but rather reflects factors that influence the competence of oocytes coming into the laboratory. The expected range is 75–90% at 40 ± 1 h post-trigger for all COC retrieved. Values outside this range could prompt a review of any changes in ovarian stimulation, triggering, or follicle aspiration practice, as changes in the proportion of MII oocytes could be a factor in changes in fertilization rates and/or embryo development. Instability in this value could indicate changes in the stimulation, resulting in a higher proportion of either immature or post-mature oocytes.

Notes:

- Since this value is expected to be stable, laboratories may choose 'reporting by exception' – only reporting on it when it falls outside the expected range.
- It should be noted that nuclear maturity does not necessarily indicate cytoplasmic maturity of the oocyte [Coticchio et al., 2012; Eppig, 1996; Sundstrom and Nilsson, 1988].
- Clinics should consider whether they should sub-divide their assessment of this Indicator based on patient demographics.
- Good communication between laboratory and clinic (cycle planning and cycle review) were cited as being vital to excellent outcomes in IVF programs [Van Voorhis et al., 2010].

Semen analysis characteristics

Sperm concentration, motility and vitality. Unless semen analyses are performed employing analytical methods as per the ESHRE SIG Andrology [Björndahl et al., 2010] or WHO5 [World Health Organization, 2010] results for sperm concentration, motility and vitality will be subject to unacceptably high uncertainty of measurement [Björndahl et al., 2016; Sanchez-Pozo et al., 2013].

Sperm morphology. Sperm morphology assessment is subjective and so is dependent on consistent training. Since the Tygerberg Strict Criteria cut-off of 4% normal forms was derived in relation to IVF success, it could be pertinent in differentiating between the need for IUI, IVF or ICSI [Menkveld, 2010; Mortimer and Menkveld, 2001]. However, the current visual evaluation of 200 or 400 spermatozoa used in the vast majority of laboratories to assess '% normal forms' has such a large uncertainty of measurement that it cannot be considered a reliable predictor for IVF success/failure for individual men [Björndahl et al., 2010; Kvist and Björndahl, 2002]. Unless determined using a more robust methodology, sperm normal forms should not be used to direct ART treatment options.

Sperm motility post-preparation

This PI is defined as the proportion of progressively motile spermatozoa in the sperm preparation for insemination, and includes only fresh normozoospermic ejaculate specimens. Sperm motility after washing should be very high, and low values would indicate problems with the preparation procedure. In this case, progressive motility

is defined as spermatozoa that are moving with net space gain of the head, and so includes hyperactivated spermatozoa. Although there was excellent agreement among the survey responses, potential weaknesses of this PI include: possible poor reliability of % motility data; non-robust classification of cases based on uncertainty of semen analysis data; variability in sperm preparation method used; and abnormal response of the sperm to the preparation method used. The reference values were competence 90% and benchmark $\geq 95\%$.

Notes:

- There is no sperm recovery rate KPI recommended because this is so heavily dependent on the processing method.
- Recommendations for IUI or IVF treatment: It was the recommendation of the Expert Panel that decisions regarding a man's suitability or need for an appropriate ART treatment modality (IUI, IVF or ICSI) should be based on number of spermatozoa and motility assessments determined in a pre-treatment 'trial preparation'. The competency threshold value was agreed as at least 90% progressive motility post-wash with a benchmark of at least 95% progressive motility.
- Recommendations for ICSI Treatment: There was no cut-off value recommended for ICSI treatment, beyond the spermatozoa ideally being alive. In this case, the best evidence of vitality is motility, the second-best evidence, in the absence of motility, is a positive hypoosmotic swelling (HOS) test [Björndahl et al., 2010], although other methodologies also exist.

ICSI damage rate

This KPI (Table 4) is defined as the proportion of oocytes that are damaged during the ICSI injection, or have degenerated by the time of fertilization assessment on Day 1. It is informative of gamete quality and/or operator skill, and excludes damage from oocyte stripping, which should be very rare. The results can be skewed by the patient mix or the stimulation protocols used, so all cycles should be included to reduce the relative impact of these variables. The Alpha survey gave similar median and mode values for each level. These values were agreed by the Expert Panel in relation to those recommended by ACE (Hughes and Association of Clinical Embryologists, 2012). The reference values for ICSI damage rate are: competence $\leq 10\%$; benchmark $\leq 5\%$.

ICSI normal fertilization rate

This KPI (Table 4) is defined as the number of fertilized oocytes on Day 1 (presence of 2PN and 2PB assessed at 17 ± 1 h post-injection) (Alpha Scientists In Reproductive Medicine and ESHRE Special Interest Group of Embryology, 2011), as a function of all MII oocytes injected. This is a common, broad, effective indicator of good laboratory practice, as it is informative of gamete quality and/or operator skill. This KPI includes ejaculated spermatozoa only (fresh or frozen) as results may be lower with surgically-retrieved spermatozoa, and excludes in-vitro matured oocytes, as well as thawed/warmed oocytes (this was dealt with in the cryopreservation consensus) (Alpha Scientists In Reproductive Medicine, 2012). Reference values were agreed by the Expert Panel: competence $\geq 65\%$; benchmark $\geq 80\%$.

Notes:

- Total ICSI failed fertilization rate: It was the consensus of the Expert Panel that complete failure to achieve normal fertilization in an ICSI cycle did not need to be a PI, but should be reported by exception, meaning that every case should be investigated. This includes only stimulated cycles, as natural cycles are expected to have only 1 oocyte.
- Poor ICSI fertilization rate: Although opinion regarding the expected incidence of cycles with ICSI fertilization rates $< 25\%$ was sought in the Alpha survey, the consensus of the Expert Panel was

to exclude this Indicator, as it did not add to the information already collected.

- Giant oocytes should not be injected due to published evidence of chromosomal abnormality (Balakier et al., 2002; Lehner et al., 2015; Rosenbusch et al., 2002).
- Regarding oocytes with smooth endoplasmic reticulum clusters (SER), more recent publications reporting outcomes suggest that the Istanbul consensus recommendation not to inject/inseminate these oocytes may need to be revisited (Mateizel et al., 2013). It was the opinion of the Expert Panel that, in the meantime, the decision to inject SER-positive oocytes should be reviewed by the clinical team on a case-by-case basis. Follow-up of results, including pregnancy outcome and babies born after insemination and transfer of the resulting embryos, should be performed.

Normal IVF fertilization rate

This KPI (Table 4) is defined as the number of fertilized oocytes on Day 1 (presence of 2PN and 2PB assessed at 17 ± 1 h post-insemination) (Alpha Scientists In Reproductive Medicine and ESHRE Special Interest Group of Embryology, 2011), as a function of all COC inseminated. IVF normal fertilization rate is an important indicator of laboratory performance, as it relies upon effective gamete handling and culture, and so is a measure of the whole in-vitro fertilization system. The reference values are: competence $\geq 60\%$; benchmark $\geq 75\%$. It should be noted that the benchmark value was determined based on a rounding of the product of the benchmark rates for MII oocytes (90%) and fertilization rate of MII oocytes (80%).

IVF polyspermy rate

This PI (Table 3) is defined as the proportion of inseminated oocytes with more than two pronuclei on Day 1 (17 ± 1 h post-insemination). It is needed to provide the information to interpret any observed variations in the normal fertilization rate. It was the consensus value that polyspermy rate should be $< 6\%$. Observed values above this rate should be reported and investigated.

1PN rate following IVF or ICSI

This PI (Table 3) is defined as the proportion of inseminated oocytes with one pronucleus on Day 1 (17 ± 1 h post-insemination). It can provide a marker of a problem in gamete handling or culture conditions and so should be low under normal conditions. It was the consensus that the 1PN rate should be $< 5\%$ for IVF cycles, and $< 3\%$ for ICSI cycles. The difference between IVF and ICSI is related to the pre-selection of oocytes prior to ICSI injection. Observed values above this rate should be reported and investigated.

Failed fertilization rate (IVF cycles)

This KPI (Table 4) is defined as the proportion of IVF cycles (excludes ICSI cycles) with no evidence of fertilization (i.e. 0 oocytes with ≥ 2 PN) on Day 1 (17 ± 1 h post-insemination). It can provide a marker of a problem in gamete quality (sperm function, oocyte activation, gamete receptors), sperm processing, or in the number of spermatozoa used for insemination. It should be low, under normal conditions. Based on the Alpha survey, and on the values recommended by ACE (Hughes and Association of Clinical Embryologists, 2012), it was the consensus that the IVF failed fertilization rate should be $< 5\%$ for stimulated cycles. Observed values above this rate should be reported and investigated.

Zygote grade (IVF cycles)

Zygote grade is an evaluation of the quality of the fertilized oocyte, conducted 17 ± 1 h post-insemination. It was the consensus that there were not enough data to recommend Indicator values for this measure.

In the discussion related to micronuclei, it was agreed that while micronuclei are abnormal, there is no evidence to confirm the identity of these dynamic manifestations.

Early cleavage rate

Early cleavage rate is the proportion of fertilized oocytes that have undergone the first round of cleavage by 26 ± 1 h post-insemination by ICSI or 28 ± 1 h post-insemination by IVF. There is evidence that early cleavage, together with other factors, can be used as an embryo selection method as it has been correlated with implantation rate [Balaban and Urman, 2003; Brezinova et al., 2009; Ciray et al., 2005; Lundqvist et al., 2001; Shoukir et al., 1997]. However, as it is not routinely calculated, it was the consensus that while this Indicator can be useful for troubleshooting purposes, there were no recommendations for expected values.

Cleavage rate

This KPI (Table 4) is defined as the proportion of zygotes that cleave to become embryos on Day 2 (44 ± 1 h post-insemination) [Alpha Scientists In Reproductive Medicine and ESHRE Special Interest Group of Embryology, 2011]. It provides an indication of the ability of the culture system to support cleavage of fertilized oocytes (i.e. with occurrence of cellular division), and of embryo viability, so a low cleavage rate could be a warning that the culture system has been impacted by an extrinsic factor. Furthermore, there is evidence that the presence of at least 1 non-cleaved embryo is predictive of reduced embryo quality for the remaining cohort [Machtinger et al., 2015]. The reference values are: competency $>95\%$; benchmark $>99\%$.

Embryo development rate

This KPI (Table 4) is defined as the proportion of cleaved embryos at the 4-cell stage on Day 2 (44 ± 1 h post-insemination) or at the 8-cell stage on Day 3 (68 ± 1 h post-insemination) per normally fertilized oocyte. This is an indicator of the ability of the culture system to support cleavage according to the expected stages, as well as providing an indication of the viability and quality of embryos. This KPI just considers the developmental stage of the embryo, regardless of grade, because developmental stage has been reported to give the highest degree of predictive power and has the advantage of being an objective measure. It was acknowledged that not all clinics consider the specific cell stages defined here, and that the culture system used can affect the kinetics of embryo development.

The reference values, based on the median and maximum results from the participants' laboratories are: Day 2, competency $\geq 50\%$ and benchmark $\geq 80\%$; and Day 3, competency $\geq 45\%$ and benchmark $\geq 70\%$.

Notes:

- It was the decision of the Expert Panel not to consider Day 4 embryo development rate.
- This KPI is most relevant to clinics that transfer embryos on Day 2 or Day 3.

Embryo and blastocyst utilization rates

These potential Indicators were defined as the number of embryos (or blastocysts) suitable for transfer or cryopreservation as a function of the number of normally fertilized (2PN) oocytes observed on Day 1. Although it was agreed that these Indicators could be of value for internal laboratory comparison, the consensus opinion was that because there are so many differences in laboratory and clinical practice, it was not practical to suggest any values for these Indicators.

Blastocyst development rate

This KPI (Table 4) is defined as the proportion of blastocysts observed at 116 ± 2 h post-insemination as a function of the number of normally fertilized oocytes. It estimates the ability of the culture system to support blastocyst formation from fertilized oocytes (i.e. with formation of ICM, trophectoderm cells and a blastocoele cavity), and provides an indication of embryo viability. It should be noted that this definition only considers blastocyst formation, with no consideration of blastocyst stage or blastocyst quality.

The reference values are: competency $\geq 40\%$; benchmark $\geq 60\%$ (Day 5). A possible additional PI might be the development of an additional 10–15% blastocysts by 140 ± 2 h post-insemination (i.e. by Day 6).

Good blastocyst development rate

This PI (Table 3) is defined as the number of good-quality blastocysts as a function of the number of normally fertilized oocytes. Blastocyst quality is as defined in the Istanbul consensus [Alpha Scientists In Reproductive Medicine and ESHRE Special Interest Group of Embryology, 2011]. This Indicator estimates the ability of the culture system to support the formation of high-grade blastocysts from fertilized oocytes (i.e. with formation of ICM, trophectoderm cells and a blastocoele cavity), and an indication of embryo viability. The reference values are: competency $\geq 30\%$; benchmark $\geq 40\%$ (Day 5). A possible additional PI might be the development of an additional 5–15% blastocysts by 140 ± 2 h post-insemination (i.e. by Day 6), depending upon the culture system.

Proportion of good blastocysts

This PI is defined as the proportion of blastocysts with a grade of 'good' or higher. There was no discussion of reference values for this Indicator, as they can be inferred from the preceding two Indicators.

Day 5 embryo transfer rate

This PI is defined as the proportion of cycles with at least one utilizable blastocyst on Day 5 relative to the presence of at least one 2PN oocyte on Day 1, to allow for the inclusion of cycles in which the decision has been made to cryopreserve all embryos. This Indicator reflects the efficiency of the whole culture system, but is only relevant for those clinics that have a blanket strategy of Day 5 transfers. It was the consensus opinion that as there are too many clinic-specific variables, including different embryo transfer policies in different centres, clinics should develop their own expectations for this Indicator, depending on when the decision to go to Day 5 transfer is made (e.g. Day 0 versus Day 3).

Implantation rate

For this consensus, this KPI (Table 4) is defined as the number of gestational sacs divided by the total number of embryos transferred, irrespective of whether a pregnancy was established [Zegers-Hochschild et al., 2009]. There was some dissent about the use of sacs, rather than fetal heartbeat – which was felt to be a more meaningful indicator of implantation rate – however sacs are used by most national/international registries. Following discussion, it was agreed to use sacs for the purpose of this consensus, but that the use of fetal heartbeat as the numerator should be revisited in the future.

Implantation rate provides an indication of the overall performance of the laboratory, so an overall low implantation rate is a serious sign of a systemic problem. Values would be expected to be lower for Day 2 and Day 3 transfers than for blastocyst transfers. In addition, results will be influenced by clinical factors (e.g. uterine receptivity) and the existence of different policies in different centres for deciding the day of embryo transfer.

Reference values for implantation rates.

Planned transfers of cleavage-stage embryos (Day 2 or Day 3): competency $\geq 25\%$; benchmark $\geq 35\%$

Blastocyst transfers: competency $\geq 35\%$; benchmark $\geq 60\%$ (the panel was divided between 55% and 60%, but agreed that 60% was an aspirational goal)

Note: These values could be affected if there are a large number of patients in the cohort who have had a large number of previous, unsuccessful cycles or significant clinical adverse factors. In addition, individual clinics may wish to further stratify their results based on patient age groups.

Live birth rate

It was the consensus that there are too many other variables to support the use of LBR as a laboratory indicator for either cleavage-stage or blastocyst-stage embryo transfers.

Successful biopsy rate

This KPI (Table 4) is defined as the proportion of biopsied and tubed/fixed samples where DNA is detected. It is a measure of the ability of embryologists to transfer the biopsied samples to test tubes, as proven by positive DNA amplification.

Based on data from the surveys and the PGD Consortium, which reported 91% diagnosis rate in 254,820 biopsies (De Rycke et al., 2015), the reference values were: competency $\geq 90\%$; benchmark $\geq 95\%$ (Table 4).

No biopsy

This Indicator was defined as the proportion of treatment cycles that had no embryos available for biopsy. It was the consensus that not having good-quality blastocysts for biopsy is not an indication of the quality of the PGD/PGS service. It is an expression of blastocyst development rates, and so no further values were developed for this Indicator.

No embryos found on warming or degenerated embryos found on warming

Nowadays, the majority of PGT cycles are based on blastocyst biopsy and cryopreservation. It was agreed that no embryos found on warming or degenerated embryos found on warming should be a KPI, as it provides a reflection of operator skill, and/or the device used. However, since not finding an embryo is a rare event, the panel was unable to estimate a competency value, as the value would be very low.

Due to greater experience with blastocyst vitrification, the rate of degeneration on warming should now be lower than that estimated in the previous cryopreservation consensus (Alpha Scientists In Reproductive Medicine, 2012). Similarly, it was the consensus that re-expansion does not differ between (warmed) biopsied and non-biopsied blastocysts.

Notwithstanding some device differences, the reference rates for blastocyst cryosurvival could now reasonably be expected to be: competency $\geq 90\%$; benchmark $\geq 99\%$ (Table 4).

Implantation rate of biopsied embryos

It was the consensus that the implantation rate for blastocysts biopsied for PGS should exceed that expected for the age-matched patient population in the same clinic. From the literature, a meta-analysis reported an improvement of 30% sustained implantation rate after the transfer of PGS-selected blastocysts relative to controls (Dahdouh et al., 2015; Scott et al., 2013).

Time-lapse imaging

Despite there being an increasing number of IVF cycles incorporating time-lapse imaging for embryo assessment and selection, the panel considered it premature to propose time-lapse related performance indicators for the IVF laboratory, due to the limited and varied data associating precise timings of human embryo development with viability or good laboratory practice. It was accepted, however, that due to the detailed morphological and kinetic information collected per embryo, time-lapse imaging may prove to be a future early warning tool for compromised culture conditions, providing a (intra-)laboratory PI, specifically if a change in mean timings for embryos to reach developmental milestones may be detected more readily and rapidly than with standard assessment methods.

A time-lapse assay of mouse embryo development linked specific morphokinetic changes to toxicity of mineral oil. This demonstrates the sensitivity of mouse embryo cleavage timings to the quality of the culture environment and the potential value of time-lapse in detecting such changes (Wolff et al., 2013). To date, no studies of this type have been performed on human embryos.

The current recommendation, therefore, was that clinics may wish to establish time-lapse KPI and benchmarks based on their own experience. It was suggested that the frequency of anomalous cleavage events, such as trichotomous mitosis (direct cleavage to three cells), which is known to be relatively common (6–8%) and associated with reduced implantation potential, could be monitored for future use as a KPI (Athayde Wirka et al., 2014; Rubio et al., 2012). In addition, it was considered that clinics using validated time-lapse algorithms for embryo selection may develop benchmarks associated with the proportion of embryos which ranked highest using morphokinetic modelling. Large-scale population studies are, however, required in order to identify which, if any, morphokinetic markers can be developed into universal and useful laboratory KPIs. It is also important to point out that morphokinetic embryo performance is believed to be highly associated with clinical and laboratory practices, and may therefore be difficult to compare between laboratories.

Acknowledgement

The consensus meeting and writing of the paper were supported by funds from ESHRE and Alpha. Alpha gratefully acknowledges the following organizations for their financial support, through the provision of unrestricted educational grants: Global Fertility Alliance, Merck KGaA, Origio, and Vitrolife. We also thank the respondents to the questionnaires for their valuable contribution to this consensus. Additionally, we would like to express our gratitude to the following societies for providing

their valuable insights on the laboratory performance indicators: BLEFCO (France), AGRBM (Germany), ICE (Ireland), SIERR (Italy), SASREG (South Africa), KED (Turkey), ACE (UK). This manuscript was prepared by Nathalie Vermeulen and Sharon Mortimer.

ARTICLE INFO

Article history:

Received 14 June 2017

Accepted 21 June 2017

Keywords:

Embryo

IVF

KPI

Oocyte

Performance indicator

Sperm

REFERENCES

- Ahlstrom, A., Westin, C., Reismer, E., Wikland, M., Hardarson, T., 2011. Trophoctoderm morphology: an important parameter for predicting live birth after single blastocyst transfer. *Hum. Reprod.* 26, 3289–3296.
- Ahlstrom, A., Westin, C., Wikland, M., Hardarson, T., 2013. Prediction of live birth in frozen-thawed single blastocyst transfer cycles by pre-freeze and post-thaw morphology. *Hum. Reprod.* 28, 1199–1209.
- Alpha Scientists In Reproductive Medicine, 2012. The Alpha consensus meeting on cryopreservation key performance indicators and benchmarks: proceedings of an expert meeting. *Reprod. Biomed. Online* 25, 146–167.
- Alpha Scientists In Reproductive Medicine, ESHRE Special Interest Group of Embryology, 2011. The Istanbul consensus workshop on embryo assessment: proceedings of an expert meeting. *Hum. Reprod.* 26, 1270–1283. and *Reprod Biomed Online* 22:1632–1646 [simultaneous publication].
- Arce, J.C., Andersen, A.N., Fernandez-Sanchez, M., Visnova, H., Bosch, E., Garcia-Velasco, J.A., Barri, P., de Sutter, P., Klein, B.M., Fauser, B.C., 2014. Ovarian response to recombinant human follicle-stimulating hormone: a randomized, antimullerian hormone-stratified, dose-response trial in women undergoing in vitro fertilization/intracytoplasmic sperm injection. *Fertil. Steril.* 102, 1633–1640, e1635.
- Athayde Wirka, K., Chen, A.A., Conaghan, J., Ivani, K., Gvakharia, M., Behr, B., Suraj, V., Tan, L., Shen, S., 2014. Atypical embryo phenotypes identified by time-lapse microscopy: high prevalence and association with embryo development. *Fertil. Steril.* 101, 1637–1648, e1631–1635.
- Baart, E.B., Martini, E., Eijkemans, M.J., Van Opstal, D., Beckers, N.G., Verhoeff, A., Macklon, N.S., Fauser, B.C., 2007. Milder ovarian stimulation for in-vitro fertilization reduces aneuploidy in the human preimplantation embryo: a randomized controlled trial. *Hum. Reprod.* 22, 980–988.
- Balaban, B., Urman, B., 2003. Embryo culture as a diagnostic tool. *Reprod. Biomed. Online* 7, 671–682.
- Balaban, B., Urman, B., 2006. Effect of oocyte morphology on embryo development and implantation. *Reprod. Biomed. Online* 12, 608–615.
- Balakier, H., Bouman, D., Sojecki, A., Librach, C., Squire, J.A., 2002. Morphological and cytogenetic analysis of human giant oocytes and giant embryos. *Hum. Reprod.* 17, 2394–2401.
- Björndahl, L., 2011. What is normal semen quality? On the use and abuse of reference limits for the interpretation of semen analysis results. *Hum. Fertil. (Camb.)* 14, 179–186.
- Björndahl, L., Barratt, C.L., Mortimer, D., Jouannet, P., 2016. 'How to count sperm properly': checklist for acceptability of studies based on human semen analysis. *Hum. Reprod.* 31, 227–232.
- Björndahl, L., Mortimer, D., Barratt, C.L., Castilla, J.A., Menkveld, R., Kvist, U., Alvarez, J.G., Haugen, T.B., 2010. A Practical Guide to Basic Laboratory Andrology. Cambridge University Press, Cambridge, UK.
- Braga, D.P., Setti, A.S., Figueira Rde, C., Machado, R.B., Iaconelli, A., Jr., Borges, E., Jr., 2013. Influence of oocyte dysmorphisms on blastocyst formation and quality. *Fertil. Steril.* 100, 748–754.
- Brezinova, J., Oborna, I., Svobodova, M., Fingerova, H., 2009. Evaluation of day one embryo quality and IVF outcome—a comparison of two scoring systems. *Reprod. Biol. Endocrinol.* 7, 9.
- Ciray, H.N., Karagenc, L., Ulug, U., Bener, F., Bahceci, M., 2005. Use of both early cleavage and day 2 mononucleation to predict embryos with high implantation potential in intracytoplasmic sperm injection cycles. *Fertil. Steril.* 84, 1411–1416.
- Cobo, A., de los Santos, M.J., Castello, D., Gamiz, P., Campos, P., Remohi, J., 2012. Outcomes of vitrified early cleavage-stage and blastocyst-stage embryos in a cryopreservation program: evaluation of 3,150 warming cycles. *Fertil. Steril.* 98, 1138–1146, e1131.
- Coetzee, K., Kruge, T.F., Lombard, C.J., 1998. Predictive value of normal sperm morphology: a structured literature review. *Hum. Reprod. Update* 4, 73–82.
- Cooper, T.G., Noonan, E., von Eckardstein, S., Auger, J., Baker, H.W., Behre, H.M., Haugen, T.B., Kruger, T., Wang, C., Mbizvo, M.T., Vogelsong, K.M., 2010. World Health Organization reference values for human semen characteristics. *Hum. Reprod. Update* 16, 231–245.
- Coticchio, G., Dal-Canto, M., Guglielmo, M.C., Mignini-Renzini, M., Fadini, R., 2012. Human oocyte maturation in vitro. *Int. J. Dev. Biol.* 56, 909–918.
- de los Santos, M.J., Arroyo, G., Busquet, A., Calderon, G., Cuadros, J., Hurtado de Mendoza, M.V., Moragas, M., Herrero, R., Ortiz, A., Pons, C., Ten, J., Vilches, M.A., Figueroa, M.J., Embryology, A.I.G.I., 2014. A multicenter prospective study to assess the effect of early cleavage on embryo quality, implantation, and live-birth rate. *Fertil. Steril.* 101, 981–987.
- Dahdouh, E.M., Balayla, J., Garcia-Velasco, J.A., 2015. Comprehensive chromosome screening improves embryo selection: a meta-analysis. *Fertil. Steril.* 104, 1503–1512.
- Davison, S.L., Bell, R., Donath, S., Montalto, J.G., Davis, S.R., 2005. Androgen levels in adult females: changes with age, menopause, and oophorectomy. *J. Clin. Endocrinol. Metab.* 90, 3847–3853.
- De Rycke, M., Belva, F., Goossens, V., Moutou, C., SenGupta, S.B., Traeger-Synodinos, J., Coonen, E., 2015. ESHRE PGD Consortium data collection XIII: cycles from January to December 2010 with pregnancy follow-up to October 2011. *Hum. Reprod.* 30, 1763–1789.
- Delvigne, A., 2009. Symposium: update on prediction and management of OHSS. *Epidemiology of OHSS. Reprod. Biomed. Online* 19, 8–13.
- Denomme, M.M., Mann, M.R., 2012. Genomic imprints as a model for the analysis of epigenetic stability during assisted reproductive technologies. *Reproduction* 144, 393–409.
- Ebner, T., Moser, M., Sommergruber, M., Tews, G., 2003. Selection based on morphological assessment of oocytes and embryos at different stages of preimplantation development: a review. *Hum. Reprod. Update* 9, 251–262.
- Ebner, T., Montag, M., Oocyte Activation Study Group, Montag, M., Van der Ven, K., Van der Ven, H., Ebner, T., Shebl, O., Oppelt, P., Hirchenhain, J., Krussel, J., Maxrath, B., Gnoth, C., Friol, K., Tigges, J., Wunsch, E., Luckhaus, J., Beerkotte, A., Weiss, D., Grunwald, K., Struller, D., Etien, C., 2015. Live birth after artificial oocyte activation using a ready-to-use ionophore: a prospective multicentre study. *Reprod. Biomed. Online* 30, 359–365.

- Ectors, F.J., Vanderzwalmen, P., Van Hoeck, J., Nijs, M., Verhaegen, G., Delvigne, A., Schoysman, R., Leroy, F., 1997. Relationship of human follicular diameter with oocyte fertilization and development after in-vitro fertilization or intracytoplasmic sperm injection. *Hum. Reprod.* 12, 2002–2005.
- Eppig, J.J., 1996. Coordination of nuclear and cytoplasmic oocyte maturation in eutherian mammals. *Reprod. Fertil. Dev.* 8, 485–489.
- ESHRE Guideline Group on Good Practice in IVF Labs, De los Santos, M.J., Apter, S., Cotichio, G., Debrock, S., Lundin, K., Plancha, C.E., Prados, F., Rienzi, L., Verheyen, G., Woodward, B., Vermeulen, N., 2016. Revised guidelines for good practice in IVF laboratories (2015). *Hum. Reprod.* 31, 685–686.
- Fancsovičs, P., Tothne, Z.G., Murber, A., Rigo, J., Jr., Urbancsek, J., 2012. Importance of cytoplasmic granularity of human oocytes in in vitro fertilization treatments. *Acta Biol. Hung.* 63, 189–201.
- Fauque, P., Jouannet, P., Lesaffre, C., Ripoche, M.A., Dandolo, L., Vaiman, D., Jammes, H., 2007. Assisted Reproductive Technology affects developmental kinetics, H19 Imprinting Control Region methylation and H19 gene expression in individual mouse embryos. *BMC Dev. Biol.* 7, 116.
- Fauque, P., Audureau, E., Leandri, R., Delaroché, L., Assouline, S., Epelboin, S., Jouannet, P., Patrat, C., 2013. Is the nuclear status of an embryo an independent factor to predict its ability to develop to term? *Fertil. Steril.* 99, 1299–1304, e1293.
- Ferraretti, A.P., Gianaroli, L., Magli, M.C., D'Angelo, A., Farfalli, V., Montanaro, N., 2004. Exogenous luteinizing hormone in controlled ovarian hyperstimulation for assisted reproduction techniques. *Fertil. Steril.* 82, 1521–1526.
- Flaherty, S.P., Payne, D., Matthews, C.D., 1998. Fertilization failures and abnormal fertilization after intracytoplasmic sperm injection. *Hum. Reprod.* 13 (Suppl. 1), 155–164.
- Garrido, N., Bellver, J., Remohi, J., Simon, C., Pellicer, A., 2011. Cumulative live-birth rates per total number of embryos needed to reach newborn in consecutive in vitro fertilization (IVF) cycles: a new approach to measuring the likelihood of IVF success. *Fertil. Steril.* 96, 40–46.
- Glujovsky, D., Farquhar, C., Quinteiro Retamar, A.M., Alvarez Sedo, C.R., Blake, D., 2016. Cleavage stage versus blastocyst stage embryo transfer in assisted reproductive technology. *Cochrane Database Syst. Rev.* (6), CD002118.
- Haaf, T., Hahn, A., Lambrecht, A., Grossmann, B., Schwaab, E., Khanaga, O., Hahn, T., Tresch, A., Schorsch, M., 2009. A high oocyte yield for intracytoplasmic sperm injection treatment is associated with an increased chromosome error rate. *Fertil. Steril.* 91, 733–738.
- Heindryckx, B., Van der Elst, J., De Sutter, P., Dhont, M., 2005. Treatment option for sperm- or oocyte-related fertilization failure: assisted oocyte activation following diagnostic heterologous ICSI. *Hum. Reprod.* 20, 2237–2241.
- Hughes, C., Association of Clinical Embryologists, 2012. Association of clinical embryologists—guidelines on good practice in clinical embryology laboratories 2012. *Hum. Fertil. (Camb.)* 15, 174–189.
- Hugues, J.N., Theron-Gerard, L., Coussieu, C., Pasquier, M., Dewailly, D., Cedrin-Durnerin, I., 2010. Assessment of theca cell function prior to controlled ovarian stimulation: the predictive value of serum basal/stimulated steroid levels. *Hum. Reprod.* 25, 228–234.
- Humaidan, P., Bungum, M., Bungum, L., Yding Andersen, C., 2004. Effects of recombinant LH supplementation in women undergoing assisted reproduction with GnRH agonist down-regulation and stimulation with recombinant FSH: an opening study. *Reprod. Biomed. Online* 8, 635–643.
- Jeppesen, J.V., Kristensen, S.G., Nielsen, M.E., Humaidan, P., Dal Canto, M., Fadini, R., Schmidt, K.T., Ernst, E., Yding Andersen, C., 2012. LH-receptor gene expression in human granulosa and cumulus cells from antral and preovulatory follicles. *J. Clin. Endocrinol. Metab.* 97, E1524–E1531.
- Joergensen, M.W., Labouriau, R., Hindkjaer, J., Stougaard, M., Kolevraa, S., Bolund, L., Agerholm, I.E., Sunde, L., 2015. The parental origin correlates with the karyotype of human embryos developing from tripronuclear zygotes. *Clin Exp Reprod Med* 42, 14–21.
- Kai, Y., Iwata, K., Iba, Y., Mio, Y., 2015. Diagnosis of abnormal human fertilization status based on pronuclear origin and/or centrosome number. *J. Assist. Reprod. Genet.* 32, 1589–1595.
- Kashir, J., Heindryckx, B., Jones, C., De Sutter, P., Parrington, J., Coward, K., 2010. Oocyte activation, phospholipase C zeta and human infertility. *Hum. Reprod. Update* 16, 690–703.
- Kohn, L.T., Corrigan, J.M., Donaldson, M.S., 2000. *To Err is Human: Building a Safer Health System*. National Academies Press.
- Kruger, T.F., Acosta, A.A., Simmons, K.F., Swanson, R.J., Matta, J.F., Oehninger, S., 1988. Predictive value of abnormal sperm morphology in in vitro fertilization. *Fertil. Steril.* 49, 112–117.
- Kvist, U., Bjorndahl, L., 2002. *Manual on Basic Semen Analysis*. Oxford University Press.
- Lehner, A., Kaszas, Z., Murber, A., Rigo, J., Jr., Urbancsek, J., Fancsovičs, P., 2015. Giant oocytes in human in vitro fertilization treatments. *Arch. Gynecol. Obstet.* 292, 697–703.
- Lemmen, J.G., Rodriguez, N.M., Andreassen, L.D., Loft, A., Ziebe, S., 2016. The total pregnancy potential per oocyte aspiration after assisted reproduction—in how many cycles are biologically competent oocytes available? *J. Assist. Reprod. Genet.* 33, 849–854.
- Levron, J., Munne, S., Willadsen, S., Rosenwaks, Z., Cohen, J., 1995. Male and female genomes associated in a single pronucleus in human zygotes. *Biol. Reprod.* 52, 653–657.
- Liu, J., Nagy, Z., Joris, H., Tournaye, H., Smitz, J., Camus, M., Devroey, P., Van Steirteghem, A., 1995. Analysis of 76 total fertilization failure cycles out of 2732 intracytoplasmic sperm injection cycles. *Hum. Reprod.* 10, 2630–2636.
- Lopes, A.S., Frederickx, V., Van Kerkhoven, G., Campo, R., Puttemans, P., Gordts, S., 2015. Survival, re-expansion and cell survival of human blastocysts following vitrification and warming using two vitrification systems. *J. Assist. Reprod. Genet.* 32, 83–90.
- Lundqvist, M., Johansson, U., Lundkvist, O., Milton, K., Westin, C., Simberg, N., 2001. Does pronuclear morphology and/or early cleavage rate predict embryo implantation potential? *Reprod. Biomed. Online* 2, 12–16.
- Machtinger, R., Bormann, C.L., Ginsburg, E.S., Racowsky, C., 2015. Is the presence of a non-cleaved embryo on day 3 associated with poorer quality of the remaining embryos in the cohort? *J. Assist. Reprod. Genet.* 32, 677–683.
- Mahutte, N.G., Arici, A., 2003. Failed fertilization: is it predictable? *Curr. Opin. Obstet. Gynecol.* 15, 211–218.
- Mateizel, I., Van Landuyt, L., Tournaye, H., Verheyen, G., 2013. Deliveries of normal healthy babies from embryos originating from oocytes showing the presence of smooth endoplasmic reticulum aggregates. *Hum. Reprod.* 28, 2111–2117.
- Mateo, S., Parriego, M., Boada, M., Vidal, F., Coroleu, B., Veiga, A., 2013. In vitro development and chromosome constitution of embryos derived from monopronucleated zygotes after intracytoplasmic sperm injection. *Fertil. Steril.* 99, 897–902, e891.
- Mayer, J.F., Jones, E.L., Dowling-Lacey, D., Nehchiri, F., Muasher, S.J., Gibbons, W.E., Oehninger, S.C., 2003. Total quality improvement in the IVF laboratory: choosing indicators of quality. *Reprod. Biomed. Online* 7, 695–699.
- Menkveld, R., 2010. Clinical significance of the low normal sperm morphology value as proposed in the fifth edition of the WHO Laboratory Manual for the Examination and Processing of Human Semen. *Asian J. Androl.* 12, 47–58.
- Menkveld, R., Holleboom, C.A., Rhemrev, J.P., 2011. Measurement and significance of sperm morphology. *Asian J. Androl.* 13, 59–68.
- Meseguer, M., Herrero, J., Tejera, A., Hilligsoe, K.M., Ramsing, N.B., Remohi, J., 2011. The use of morphokinetics as a predictor of embryo implantation. *Hum. Reprod.* 26, 2658–2671.
- Montag, M., van der Ven, H., German Pronuclear Morphology Study Group, 2001. Evaluation of pronuclear morphology as the only

- selection criterion for further embryo culture and transfer: results of a prospective multicentre study. *Hum. Reprod.* 16, 2384–2389.
- Mortimer, D., Menkveld, R., 2001. Sperm morphology assessment—historical perspectives and current opinions. *J. Androl.* 22, 192–205.
- Mortimer, S.T., Mortimer, D., 2015. *Quality and Risk Management in the IVF Laboratory*. Cambridge University Press.
- Nagy, Z.P., Liu, J., Joris, H., Verheyen, G., Tournaye, H., Camus, M., Derde, M.C., Devroey, P., Van Steirteghem, A.C., 1995. The result of intracytoplasmic sperm injection is not related to any of the three basic sperm parameters. *Hum. Reprod.* 10, 1123–1129.
- Nagy, Z.P., Janssenswillen, C., Janssens, R., De Vos, A., Staessen, C., Van de Velde, H., Van Steirteghem, A.C., 1998. Timing of oocyte activation, pronucleus formation and cleavage in humans after intracytoplasmic sperm injection (ICSI) with testicular spermatozoa and after ICSI or in-vitro fertilization on sibling oocytes with ejaculated spermatozoa. *Hum. Reprod.* 13, 1606–1612.
- Nagy, Z.P., Jones-Colon, S., Roos, P., Botros, L., Greco, E., Dasig, J., Behr, B., 2009. Metabolic assessment of oocyte viability. *Reprod. Biomed. Online* 18, 219–225.
- Nel-Themaat, L., Nagy, Z.P., 2011. A review of the promises and pitfalls of oocyte and embryo metabolomics. *Placenta* 32 (Suppl. 3), S257–S263.
- Nogueira, D., Friedler, S., Schachter, M., Raziell, A., Ron-El, R., Smitz, J., 2006. Oocyte maturity and preimplantation development in relation to follicle diameter in gonadotropin-releasing hormone agonist or antagonist treatments. *Fertil. Steril.* 85, 578–583.
- Paternot, G., Wetzels, A.M., Thonon, F., Vansteenbrugge, A., Willemsen, D., Devroey, J., Debrock, S., D'Hooghe, T.M., Spiessens, C., 2011. Intra- and interobserver analysis in the morphological assessment of early stage embryos during an IVF procedure: a multicentre study. *Reprod. Biol. Endocrinol.* 9, 127.
- Patrizio, P., Sakkas, D., 2009. From oocyte to baby: a clinical evaluation of the biological efficiency of in vitro fertilization. *Fertil. Steril.* 91, 1061–1066.
- Patrizio, P., Fragouli, E., Bianchi, V., Borini, A., Wells, D., 2007. Molecular methods for selection of the ideal oocyte. *Reprod. Biomed. Online* 15, 346–353.
- Piltonen, T., Koivunen, R., Ruokonen, A., Tapanainen, J.S., 2003. Ovarian age-related responsiveness to human chorionic gonadotropin. *J. Clin. Endocrinol. Metab.* 88, 3327–3332.
- Rhenman, A., Berglund, L., Brodin, T., Olovsson, M., Milton, K., Hadziiosmanovic, N., Holte, J., 2015. Which set of embryo variables is most predictive for live birth? A prospective study in 6252 single embryo transfers to construct an embryo score for the ranking and selection of embryos. *Hum. Reprod.* 30, 28–36.
- Rienzi, L., Vajta, G., Ubaldi, F., 2011. Predictive value of oocyte morphology in human IVF: a systematic review of the literature. *Hum. Reprod. Update* 17, 34–45.
- Rizk, B., 2009. Symposium: update on prediction and management of OHSS. *Genetics of ovarian hyperstimulation syndrome*. *Reprod. Biomed. Online* 19, 14–27.
- Rosenbusch, B., Schneider, M., Glaser, B., Brucker, C., 2002. Cytogenetic analysis of giant oocytes and zygotes to assess their relevance for the development of digynic triploidy. *Hum. Reprod.* 17, 2388–2393.
- Rubino, P., Vigano, P., Luddi, A., Piomboni, P., 2016. The ICSI procedure from past to future: a systematic review of the more controversial aspects. *Hum. Reprod. Update* 22, 194–227.
- Rubio, I., Kuhlmann, R., Agerholm, I., Kirk, J., Herrero, J., Escriba, M.J., Bellver, J., Meseguer, M., 2012. Limited implantation success of direct-cleaved human zygotes: a time-lapse study. *Fertil. Steril.* 98, 1458–1463.
- Russell, D.L., Salustri, A., 2006. Extracellular matrix of the cumulus-oocyte complex. *Semin. Reprod. Med.* 24, 217–227.
- Saenz-de-Juano, M.D., Billooye, K., Smitz, J., Anckaert, E., 2016. The loss of imprinted DNA methylation in mouse blastocysts is inflicted to a similar extent by in vitro follicle culture and ovulation induction. *Mol. Hum. Reprod.* 22, 427–441.
- Salinas, M., Lopez-Garrigos, M., Gutierrez, M., Lugo, J., Sirvent, J.V., Uris, J., 2010. Achieving continuous improvement in laboratory organization through performance measurements: a seven-year experience. *Clin. Chem. Lab. Med.* 48, 57–61.
- Sanchez-Pozo, M.C., Mendiola, J., Serrano, M., Mozas, J., Bjorndahl, L., Menkveld, R., Lewis, S.E., Mortimer, D., Jorgensen, N., Barratt, C.L., Fernandez, M.F., Castilla, J.A., Special Interest Group in Andrology of the European Society of Human Reproduction and Embryology, 2013. Proposal of guidelines for the appraisal of SEMen QUALity studies (SEMQUA). *Hum. Reprod.* 28, 10–21.
- Scott, R.T., Hofmann, G.E., Muasher, S.J., Acosta, A.A., Kreiner, D.K., Rosenwaks, Z., 1989. Correlation of follicular diameter with oocyte recovery and maturity at the time of transvaginal follicular aspiration. *J. In Vitro Fert. Embryo Transf.* 6, 73–75.
- Scott, R.T., Jr., Upham, K.M., Forman, E.J., Hong, K.H., Scott, K.L., Taylor, D., Tao, X., Treff, N.R., 2013. Blastocyst biopsy with comprehensive chromosome screening and fresh embryo transfer significantly increases in vitro fertilization implantation and delivery rates: a randomized controlled trial. *Fertil. Steril.* 100, 697–703.
- Shahangian, S., Snyder, S.R., 2009. Laboratory medicine quality indicators: a review of the literature. *Am. J. Clin. Pathol.* 131, 418–431.
- Shoukir, Y., Campana, A., Farley, T., Sakkas, D., 1997. Early cleavage of in-vitro fertilized human embryos to the 2-cell stage: a novel indicator of embryo quality and viability. *Hum. Reprod.* 12, 1531–1536.
- Soares, S.R., Rubio, C., Rodrigo, L., Simón, C., Remohí, J., Pellicer, A., 2003. High frequency of chromosomal abnormalities in embryos obtained from oocyte donation cycles. *Fertil. Steril.* 80, 656–657.
- Spencer, J.B., Klein, M., Kumar, A., Azziz, R., 2007. The age-associated decline of androgens in reproductive age and menopausal Black and White women. *J. Clin. Endocrinol. Metab.* 92, 4730–4733.
- Staessen, C., Van Steirteghem, A.C., 1997. The chromosomal constitution of embryos developing from abnormally fertilized oocytes after intracytoplasmic sperm injection and conventional in-vitro fertilization. *Hum. Reprod.* 12, 321–327.
- Sugimura, S., Ritter, L.J., Rose, R.D., Thompson, J.G., Smitz, J., Mottershead, D.G., Gilchrist, R.B., 2015. Promotion of EGF receptor signaling improves the quality of low developmental competence oocytes. *Dev. Biol.* 403, 139–149.
- Sultan, K.M., Munne, S., Palermo, G.D., Alikani, M., Cohen, J., 1995. Chromosomal status of uni-pronuclear human zygotes following in-vitro fertilization and intracytoplasmic sperm injection. *Hum. Reprod.* 10, 132–136.
- Sundstrom, P., Nilsson, B.O., 1988. Meiotic and cytoplasmic maturation of oocytes collected in stimulated cycles is asynchronous. *Hum. Reprod.* 3, 613–619.
- Sundstrom, P., Saldeen, P., 2008. Early embryo cleavage and day 2 mononucleation after intracytoplasmic sperm injection for predicting embryo implantation potential in single embryo transfer cycles. *Fertil. Steril.* 89, 475–477.
- Thurin, A., Hardarson, T., Hausken, J., Jablonowska, B., Lundin, K., Pinborg, A., Bergh, C., 2005. Predictors of ongoing implantation in IVF in a good prognosis group of patients. *Hum. Reprod.* 20, 1876–1880.
- Van den Abbeel, E., Balaban, B., Ziebe, S., Lundin, K., Cuesta, M.J., Klein, B.M., Helmgard, L., Arce, J.C., 2013. Association between blastocyst morphology and outcome of single-blastocyst transfer. *Reprod. Biomed. Online* 27, 353–361.
- Van Royen, E., Mangelschots, K., De Neubourg, D., Valkenburg, M., Van de Meerssche, M., Ryckaert, G., Eestermans, W., Gerris, J., 1999. Characterization of a top quality embryo, a step towards single-embryo transfer. *Hum. Reprod.* 14, 2345–2349.
- Van Voorhis, B.J., Thomas, M., Surrey, E.S., Sparks, A., 2010. What do consistently high-performing in vitro fertilization programs in the U.S. do? *Fertil. Steril.* 94, 1346–1349.
- Wang, S.S., Sun, H.X., 2014. Blastocyst transfer ameliorates live birth rate compared with cleavage-stage embryos transfer in fresh in

- vitro fertilization or intracytoplasmic sperm injection cycles: reviews and meta-analysis. *Yonsei Med. J.* 55, 815–825.
- Wang, W., Zhang, X.H., Wang, W.H., Liu, Y.L., Zhao, L.H., Xue, S.L., Yang, K.H., 2011. The time interval between hCG priming and oocyte retrieval in ART program: a meta-analysis. *J. Assist. Reprod. Genet.* 28, 901–910.
- Wang, W.H., Day, B.N., Wu, G.M., 2003. How does polyspermy happen in mammalian oocytes? *Microsc. Res. Tech.* 61, 335–341.
- Wolff, H.S., Fredrickson, J.R., Walker, D.L., Morbeck, D.E., 2013. Advances in quality control: mouse embryo morphokinetics are sensitive markers of in vitro stress. *Hum. Reprod.* 28, 1776–1782.
- World Health Organization, 2010. WHO Laboratory Manual for the Examination and Processing of Human Semen. p. 287.
- Yanagida, K., 2004. Complete fertilization failure in ICSI. *Hum. Cell* 17, 187–193.
- Yang, W.J., Hwang, Y.C., Lin, C.S., Hwu, Y.M., Lee, R.K., Hsiao, S.Y., 2015. Embryonic early-cleavage rate is decreased with aging in GnRH agonist but not in antagonist protocols. *J. Assist. Reprod. Genet.* 32, 789–795.
- Zamah, A.M., Hsieh, M., Chen, J., Vigne, J.L., Rosen, M.P., Cedars, M.I., Conti, M., 2010. Human oocyte maturation is dependent on LH-stimulated accumulation of the epidermal growth factor-like growth factor, amphiregulin. *Hum. Reprod.* 25, 2569–2578.
- Zegers-Hochschild, F., Adamson, G.D., de Mouzon, J., Ishihara, O., Mansour, R., Nygren, K., Sullivan, E., van der Poel, S., International Committee for Monitoring Assisted Reproductive Technology, World Health Organization, 2009. The International Committee for Monitoring Assisted Reproductive Technology (ICMART) and the World Health Organization (WHO) Revised Glossary on ART Terminology, 2009. *Hum. Reprod.* 24, 2683–2687.